arXiv:hep-ph/0306031v1 3 Jun 2003

The Mixing Angle of the Lightest Scalar Nonet

José A. Oller

Departamento de Física. Universidad de Murcia.
E-30071, Murcia. Spain.
oller@um.es

Abstract

We show that the κ, $a_0(980)$, σ and the $f_0(980)$ resonances constitute the lightest scalar nonet in three different and complementary ways. First, by establishing the continuous movement of the poles from the physical to a SU(3) limit. Second, by performing an analysis of the couplings of the scalar mesons to pairs of pseudoscalars and third, by analysing the couplings of the scalars with meson-meson SU(3) scattering eigenstates. Every of the last two methods agree that the mixing angle between the singlet and the octet $I=0$ states is $\theta = 19^o \pm 5$ degrees, so that the σ is mainly the singlet and the $f_0(980)$ the isosinglet octet state.

1 Introduction

Just after the experimental discovery of the $f_0(980)$ [1], a few years after the one of the $a_0(980)$ [2], Morgan in ref.[3] made a first attempt to establish the states belonging to the $J^{PC} = 0^{++}$ lightest scalar nonet and to calculate the mixing angle. He established, as a tentative solution, that the $f_0(980)$, $a_0(980)$ and a $\kappa(\backsim 1200)$ together with an extra $f_0(\backsim 1100)$ would form this scalar nonet with $\cos^2\theta = 0.13$, being θ the mixing angle. The $a_0(980)$ with $I = 1$ and the κ with $I = 1/2$ are supposed to remain as pure octet states in this and in any other $SU(3)$ analysis that we refer here. In our notation, $\cos\theta$ is the projection of the $f_0(980)$ on the isosinglet octet state. The result of Morgan's analysis is far from the ideal mixing, the one found in the vector and tensor multiplets [4, 5, 6, 7], where $\cos^2\theta = 2/3 \simeq 0.67$, so that the state coupling more strongly with strangeness is mainly an octet. Notice that in the $\bar{q}q$ picture for mesons, the ideal mixing implies that the heavier state is $\bar{s}s$ while the lighter is $(\bar{u}u + \bar{d}d)/\sqrt{2}$, the standard example being the ϕ and ω resonances. Later on, Jaffe [8] developed his model based on the MIT Bag Model [9] for multiquark hadrons as $\bar{q}^2q^2$ states. In this calculation a 0^{++} nonet of scalar resonances emerges with rather light masses due to strong attractive gluon-magnetic interactions. The nonet is made up by a $f_0(\backsim 700)$, $f_0(980)$, $a_0(980)$, the latter two with predicted masses of 1100 MeV, and a $\kappa(900)$. In this model a natural explanation of the approximate degeneracy of the $f_0(980)$ and $a_0(980)$ is obtained in terms of just their quark content, something that badly fails when these resonances are supposed to be $\bar{q}q$ states. Here, one ends with a dual situation with respect to the just referred ideal mixing, so that $\cos^2\theta = 1/3 = 0.33$. Another pioneering work on the scalar nonet is that of Scadron [10], where making use of a quark model, he establishes as well that the lightest scalar nonet comprises the $\sigma(750)$, $\kappa(800)$, $f_0(980)$ and $a_0(980)$ so that $\cos^2\theta = 0.39$. More recently, Napsuciale [11], in the $U(3) \times U(3)$ linear sigma model with $U_A(1)$ symmetry breaking [14, 15, 16], predicted again that the lightest scalar nonet consists of the $\sigma(\backsim 500)$, $f_0(980)$, $\kappa(\backsim 900)$ and the $a_0(980)$. In this calculation the properties of the pseudoscalar nonet 0^{-+} are fitted and those of the scalar nonet are predicted. In this way, Napsuciale concludes that $\cos^2\theta = 0.87$. It is worth mentioning that both Napsuciale and Scadron use the so called strange and non-strange basis, the one that results from ideal mixing, and they obtain the same absolute value for the mixing angle in this basis, $|\phi| \simeq 15^0$, but differ in the sign, so that $\theta = 35.3 + \phi$ is also different. Refs.[12, 13] also employ the same model of ref.[11] but with rather different conclusions due to the strong sensitivity of the results of this model in terms of the pseudoscalar experimental input, see ref.[11] for a detailed discussion on this issue. Finally, Black et al. in ref.[17], from an analysis of $\pi\pi$ and $K\pi$ scattering from chiral invariant Lagrangians including bare resonances, also consider the same resonances for the lightest scalar nonet with $\cos^2\theta = 0.63$. There are several other groups that also agree that the members of the lightest scalar resonances are the ones listed above, that is, $f_0(600)$ or σ, $f_0(980)$, $\kappa(700-900)$ and $a_0(980)$ [18, 19, 20, 21, 22, 23, 24] and an accord is nowadays rising that the lightest $\bar{q}q$ nonet, with rather standard properties, mixed with a glueball is made up by more massive resonances around the 1.4 GeV region, the $K_0^*(1430)$, $a_0(1450)$, $f_0(1370)$, $f_0(1500)$ and $f_0(1710)$ [25, 26]. From the experimental point of view, new and precise experimental data has also helped to clarify to large extend the status of the lightest scalar nonet. On the one hand, we have the ϕ radiative decays to $\gamma\pi\pi$ and $\gamma\pi\eta$ [27, 28, 29, 30, 31], which rule out simple $\bar{q}q$ models for the $f_0(980)$ and $a_0(980)$ resonances [32, 25], and on the other hand we have the D^+ decays, where the σ and κ resonances are clearly seen [33, 34] with masses around 0.5 and 0.8 GeV and large widths, between 0.3–0.4 GeV.

In this work, we take the strong S-wave T-matrices from refs.[22, 35] and analyse their spectroscopic content regarding the lightest scalar mesons. It is worth stressing that these T-matrices have been by now checked in a wide range of physical processes, namely, pure strong meson-meson scattering [22, 35], $\gamma\gamma$ fusion into two mesons [36], J/Ψ decays [37] and ϕ radiative decays [38, 28]. In ref.[22] was already established that the σ, $f_0(980)$, κ and $a_0(980)$ belong to the same scalar nonet but no attempt was made to study the mixing between the singlet and the $I = 0$ state of the octet making up the σ, $f_0(980)$ resonances and to perform a SU(3) analysis of the couplings constants of the members of this lightest scalar nonet as similarly performed in the vector or tensor nonets [4, 5, 6]. Here we aim to fill up this gap. We perform two SU(3) analyses of the couplings constants of the scalar resonances by considering in the first analysis the couplings to two pseudoscalars isospin states and in the second one, the couplings to SU(3) eigenstates. The two methods have different sensitivity to the SU(3) breaking, follow very different numerical analyses, and their agreement is taken as a guarantee of the consistency of our results. This is performed in section 2 where we also show the continuity of the movement of the poles from the physical situation to a SU(3) symmetric one, where the poles corresponding to the aforementioned scalar resonances bunch in a degenerate octet of resonances and in a lighter singlet pole. Some conclusions are presented in section 3.

2 SU(3)-Analyses

In table 1 we show the physical pole positions in the second Riemann sheet[#1] from the T-matrices of refs.[22, 35]. These T-matrices are obtained once the chiral expansion is resummed to fulfill the unitarity cut to all orders. The resummation is performed so that the chiral perturbative series [39, 40] is recovered algebraically at low energies when the chiral power counting holds. In addition to refs.[22, 35], a general discussion of the method is given in ref.[41]. In fig.1 we show by the solid lines the full results of ref.[22], corresponding to the fit in eq.(73) of the latter reference. In this reference, the $\pi\pi$, $K\bar{K}$ and $\eta_8\eta_8$ states were included in the $I = 0$ channel, the $K\pi$ and $K\eta_8$ in the $I = 1/2$ one and the $\pi\eta_8$ and $K\bar{K}$ for $I = 1$.

In ref.[35] a three-momentum cut-off was used to calculate the unitarity two meson diagram, $g(s)$, but now we have calculated it through a dispersion relation so that we have a subtraction constant as free parameter for each channel. In the SU(3) limit, as shown in ref.[42], all the subtraction constants must be equal. Taken this to be case, we fit the common subtraction constant for all the isospin channels with the preferred value,

$$a(\mu) = -1.23, \text{ for } \mu = M_\rho \ , \tag{2.1}$$

where μ is the subtraction scale. The formalism is invariant under changes of the subtraction scale μ [22] such that $a(\mu') = a(\mu) + \log \mu'^2/\mu^2$. In this fit we have also included the $\eta_8\eta_8$ state for $I = 0$ as in ref.[22]. The results are given by the long-dashed lines in fig.1. Let us notice

---

[#1] The sheets of the complex s-plane are defined with respect to the branch cuts of the channel three-momenta q_i. The first or physical sheet corresponds to $\mathrm{Im}q_i \geq 0$ for all the channels, and the second sheet corresponds to changing the sign of the momenta of only the lightest channel, $\mathrm{Im}q_1 \leq 0$, while for the rest of channels $\mathrm{Im}q_i \geq 0$, $i \geq 2$.

that the differences with respect to the full results of ref.[22], solid lines, come from the fact that in this fit we have removed the set of preexisting resonances of ref.[22]. This comprises an octet of mass around 1.4 GeV and a singlet of mass around 1 GeV. The role played by the heavier octet is apparent when considering the figures in the region above 1.2 GeV. We see that the main differences for lower energies correspond to the $I = 0$ inelasticity, third panel from top to bottom and left to right.

It is also interesting to remark that by performing a comparison between the $g(s)$ function calculated with a cut-off or with the dispersive integral [20], we can then obtain an expression for the subtraction constant in terms of a natural-sized cut-off. This expression corresponds to:

$$a(\mu) = -\log \frac{4Q^2}{\mu^2} \ .$$

Thus, for $Q \simeq M_\rho \simeq 4\pi f_\pi$ GeV, we have $a(M_\rho)$ in the interval $[-2, -1.4]$, where f_π is the weak pion decay constant. Indeed, the value given above for $a(M_\rho)$ in eq.(2.1) corresponds to $Q = 712$ MeV, a rather natural value for Q. This is important in order to conclude that the physics encoded in this subtraction constant just corresponds to re-scattering between the mesons, because if the subtraction constant required unnatural values of the cut-off then one should conclude that some intrinsic dynamics is playing a role besides pure meson-meson self-interactions.

In addition, we also consider the situation where we do not include either the $\eta_8\eta_8$ state, like in ref.[35], but allowing for different values of the subtraction constants. Performing the fit we finally have:

$$a_{\pi\pi} = -1.1,\ a_{K\bar{K}} = -1.6,\ a_{K\pi} = -0.8,\ a_{K\eta} = -0.8\ a_{\pi\eta} = -0.5\ . \tag{2.2}$$

The resulting curves are the short-dashed ones of fig.1.

We take these three T-matrices as reflecting the uncertainty in our approach for the coupling constants and pole positions presented in table 1. The point is that the $\eta_8\eta_8$ channel plays some role above 1 GeV, see for example the long-dashed line in the $(1-\eta_0^2)/4$ panel, but on the other hand one can also reproduce the data without this channel, short-dashed lines. Indeed, as already noted in ref.[22], the inelasticity for the $I = 0$ S-wave partial wave, η_0, cannot be appropriately reproduced when the $\eta_8\eta_8$ channel is taken into account unless a bare singlet pole around 1 GeV is included [22], compare the long-dashed and solid lines in fig.1. Thus, although the preexisting resonances can also be avoided for energies lower than 1.2 GeV, they certainly give some contribution particularly in connection with he $\eta_8\eta_8$ channel. We have then considered the three situations presented above, where the $\eta_8\eta_8$ as well as the preexisting resonances are included [22] (solid lines), where the preexisting resonances are removed but the $\eta_8\eta_8$ is kept (long-dashed lines) and finally when both the preexisting resonances and the $\eta_8\eta_8$ channel are removed (short-dashed lines). On the other hand, it is apparent from fig.1 that the bare poles are not negligible above 1.2 GeV.



Figure 1: Fits to scattering data. The solid lines is the full approach of ref.[22]. The long-dashed lines correspond to the fit given in eq.(2.1) where the bare poles are removed and a common subtraction constant is used for all the channels. The short-dashed lines correspond to remove both the bare poles and the $\eta_8\eta_8$ channel but allowing for different values of the subtraction constants, eq.(2.2). From top to bottom and left to right, the first three panels refer to the isosinglet S-wave sector: Elastic $\pi\pi$ phase shifts, the inelastic $\pi\pi \to K\bar{K}$ phase shifts and $(1-\eta_0^2)/4$ with η_0 the inelasticity, in order. The fourth panel is the isodoblet S-wave $K\pi$ elastic phase shifts and the last one is an event distribution around the isovector $a_0(980)$ resonance.

We now perform a smooth transition from the physical limit to a SU(3) symmetric one through a continuous parameter $\lambda \in [0,1]$, such that:

$$m_\pi(\lambda) \;=\; m_\pi + \lambda(m_0 - m_\pi)\;,$$

$$
\begin{aligned}
m_K(\lambda) &= m_K + \lambda(m_0 - m_K) \ , \\
m_\eta(\lambda) &= m_\eta + \lambda(m_0 - m_\eta) \ , \\
f_K(\lambda) &= f_K + \lambda(f_\pi - f_K) \ , \\
f_\eta(\lambda) &= f_\eta + \lambda(f_\pi - f_\eta) \ .
\end{aligned}
\tag{2.3}
$$

So, for $\lambda = 0$ one has the physical limit while for $\lambda = 1$ one ends with a SU(3) symmetric point. The interesting point for us is to note the continuous movement of the pole locations of the κ, $a_0(980)$, σ and $f_0(980)$ from their physical positions ($\lambda = 0$) to a degenerate octet plus a singlet as shown in fig.2, for a common SU(3) mass $m_0 = 350$ MeV and a step in λ of 0.1. This clearly shows that an octet and a singlet of resonances in the SU(3) limit evolve and give rise to the physical poles of the lightest scalars so that these resonances must be considered as the lightest nonet of scalar resonances. Thus, the κ and $a_0(980)$ are pure octet states while the σ and $f_0(980)$ are a mixture of the singlet and octet $I = 0$ states.



Figure 2: Continuous movement of the octet and singlet poles from a SU(3) symmetric point to the physical limit. The different isospin channels are indicated in the figure and the step in the λ parameter is 0.1.

We now perform a standard SU(3) analysis of the coupling constants of the lightest scalar resonances to pairs of pseudoscalars within the aforementioned SU(3) assignment. The coupling constants are obtained by determining the residues at the pole positions of the resonances through the expression:

$$
T_{ij} = \lim_{s \to s_R} - \frac{g_i g_j}{s - s_R} \ , \tag{2.4}
$$

where s_R is the corresponding pole position and the subscripts i and j indicate the two pseudoscalar channels.

Then, for the different options taken above to fit data we show in table 1 the modules of the couplings since the phases can be contaminated from background contributions [43].

Resonance	Full ref.[22] GeV	No bare poles eq.(2.1) GeV	No bare poles eq.(2.2) GeV	Average GeV
σ	$0.445 - i\,0.220$	$0.443 - i\,0.213$	$0.442 - i\,0.214$	$(0.443 \pm 0.002) - i\,(0.216 \pm 0.004)$
	$\lvert g_{\pi\pi}\rvert = 3.01$	$\lvert g_{\pi\pi}\rvert = 2.94$	$\lvert g_{\pi\pi}\rvert = 2.95$	$\lvert g_{\pi\pi}\rvert = 2.97 \pm 0.04$
	$\lvert g_{K\bar{K}}\rvert = 1.09$	$\lvert g_{K\bar{K}}\rvert = 1.30$	$\lvert g_{K\bar{K}}\rvert = 1.34$	$\lvert g_{K\bar{K}}\rvert = 1.24 \pm 0.13$
	$\lvert g_{\eta_8\eta_8}\rvert = 0.09$	$\lvert g_{\eta_8\eta_8}\rvert = 0.04$		$\lvert g_{\eta_8\eta_8}\rvert = 0.07 \pm 0.04$
$f_0(980)$	$0.988 - i\,0.014$	$0.983 - i\,0.007$	$0.987 - i\,0.011$	$(0.986 \pm 0.003) - i\,(0.011 \pm 0.004)$
	$\lvert g_{\pi\pi}\rvert = 1.33$	$\lvert g_{\pi\pi}\rvert = 0.89$	$\lvert g_{\pi\pi}\rvert = 1.18$	$\lvert g_{\pi\pi}\rvert = 1.1 \pm 0.2$
	$\lvert g_{K\bar{K}}\rvert = 3.63$	$\lvert g_{K\bar{K}}\rvert = 3.59$	$\lvert g_{K\bar{K}}\rvert = 3.83$	$\lvert g_{K\bar{K}}\rvert = 3.68 \pm 0.13$
	$\lvert g_{\eta_8\eta_8}\rvert = 2.85$	$\lvert g_{\eta_8\eta_8}\rvert = 2.61$		$\lvert g_{\eta_8\eta_8}\rvert = 2.73 \pm 0.17$
$a_0(980)$	$1.055 - i\,0.025$	$1.032 - i\,0.042$	$1.030 - i\,0.086$	$(1.039 \pm 0.014)) - i\,(0.05 \pm 0.03)$
	$\lvert g_{\pi\eta_8}\rvert = 3.88$	$\lvert g_{\pi\eta_8}\rvert = 3.67$	$\lvert g_{\pi\eta_8}\rvert = 4.08$	$\lvert g_{\pi\eta_8}\rvert = 3.9 \pm 0.2$
	$\lvert g_{K\bar{K}}\rvert = 5.50$	$\lvert g_{K\bar{K}}\rvert = 5.39$	$\lvert g_{K\bar{K}}\rvert = 5.60$	$\lvert g_{K\bar{K}}\rvert = 5.50 \pm 0.11$
κ	$0.784 - i\,0.327$	$0.804 - i\,0.285$	$0.774 - i\,0.338$	$(0.787 \pm 0.015) - i\,(0.32 \pm 0.03)$
	$\lvert g_{K\pi}\rvert = 5.02$	$\lvert g_{K\pi}\rvert = 4.93$	$\lvert g_{K\pi}\rvert = 4.89$	$\lvert g_{K\pi}\rvert = 4.94 \pm 0.07$
	$\lvert g_{K\eta_8}\rvert = 3.10$	$\lvert g_{K\eta_8}\rvert = 2.96$	$\lvert g_{K\eta_8}\rvert = 3.00$	$\lvert g_{K\eta_8}\rvert = 3.02 \pm 0.07$

Table 1: Coupling constants for the scalar resonances for the solutions of ref.[22] and those corresponding to eqs.(2.1) and (2.2). Let us remember that the $I = 0$ states $|\pi\pi\rangle_0$ and $|\eta_8\eta_8\rangle_0$ of eq.(2.6) are normalized to 1/2 and not to 1 so as to take care of the invariance of these states under the exchange of the two π or η_8. In the fifth column we show the average of the values in the three previous columns.

Let us consider first the $a_0(980)$ and κ resonances that do not have singlet contribution and should be considered as pure octet resonances. Note that although one can also have an anti-symmetric octet representation, 8_a, from the tensorial product $8 \otimes 8$, is not allowed that a pair of pseudoscalars in the SU(3) limit stay in S-wave in this representation because of Bose-Einstein statistics. For the reader convenience, we write the scattering states with definite SU(3) eigenvalues for the symmetric octet, 8_s, and the singlet, 1:

$$\begin{aligned}
|8_s, I=1, I_3=0\rangle &= \frac{1}{\sqrt{5}}|\pi^0\eta_8\rangle + \frac{1}{\sqrt{5}}|\eta_8\pi^0\rangle + \sqrt{\frac{3}{20}}(|K^+K^-\rangle - |K^0\bar{K}^0\rangle) + \sqrt{\frac{3}{20}}(|K^-K^+\rangle - |\bar{K}^0K^0\rangle) \\
&= \frac{1}{\sqrt{5}}|\pi\eta_8\rangle + \frac{1}{\sqrt{5}}|\eta_8\pi\rangle - \sqrt{\frac{3}{10}}|K\bar{K}\rangle_1 - \sqrt{\frac{3}{10}}|\bar{K}K\rangle_1 \,, \\
|8_s, I=\frac{1}{2}, I_3=\frac{1}{2}\rangle &= \sqrt{\frac{3}{20}}(|\pi^0K^+\rangle + |K^+\pi^0\rangle) + \sqrt{\frac{3}{10}}(|\pi^+K^0\rangle + |K^0\pi^+\rangle) - \frac{1}{\sqrt{20}}(|K^+\eta_8\rangle + |\eta_8K^+\rangle) \\
&= -\frac{3}{\sqrt{20}}|\pi K\rangle_{1/2} + \frac{3}{\sqrt{20}}|K\pi\rangle_{1/2} - \frac{1}{\sqrt{20}}|K\eta_8\rangle - \frac{1}{\sqrt{20}}|\eta_8K\rangle \,,
\end{aligned}$$

$$
\begin{aligned}
|8_s, I=0, I_3=0\rangle &= \frac{1}{\sqrt{5}}(|\pi^0\pi^0\rangle + |\pi^+\pi^-\rangle + |\pi^-\pi^+\rangle) - \frac{1}{\sqrt{20}}(|K^+K^-\rangle + |K^-K^+\rangle \\
&+ |K^0\bar{K}^0\rangle + |\bar{K}^0K^0\rangle) - \frac{1}{\sqrt{5}}|\eta_8\eta_8\rangle \\
&= -\sqrt{\frac{6}{5}}|\pi\pi\rangle_0 + \frac{1}{\sqrt{10}}|K\bar{K}\rangle_0 - \frac{1}{\sqrt{10}}|\bar{K}K\rangle_0 - \sqrt{\frac{2}{5}}|\eta_8\eta_8\rangle_0 \ , \\
|1, I=0, I_3=0\rangle &= \frac{1}{\sqrt{8}}(\pi^0\pi^0 + \pi^+\pi^- + \pi^-\pi^+ + \eta_8\eta_8 + K^+K^- + K^-K^+ + K^0\bar{K}^0 + \bar{K}^0K^0) \\
&= -\frac{\sqrt{3}}{2}|\pi\pi\rangle_0 + \frac{1}{2}|\eta_8\eta_8\rangle_0 - \frac{1}{2}|K\bar{K}\rangle_0 + \frac{1}{2}|\bar{K}K\rangle_0 \ . \qquad (2.5)
\end{aligned}
$$

Above we have also shown the decomposition of the octet or singlet states in isospin states in addition to its decomposition in two pseudoscalar pairs. Furthermore, since the $|\pi\pi\rangle_0$ and $|\eta_8\eta_8\rangle$ are symmetric states under the exchange of the two pseudoscalars, we have added an extra $1/\sqrt{2}$ in the expression for the isospin states, so that:

$$
\begin{aligned}
|\pi\pi\rangle_0 &= -\frac{1}{\sqrt{6}}|\pi^0\pi^0 + \pi^+\pi^- + \pi^-\pi^+\rangle \ , \\
|\eta_8\eta_8\rangle_0 &= \frac{1}{\sqrt{2}}|\eta_8\eta_8\rangle \ . \qquad (2.6)
\end{aligned}
$$

From this decomposition and identifying the $a_0(980)$ as the $|8_s, I=1, I_3=0\rangle$ state and the κ^+ as the $|8_s, I=1/2, I_3=1/2\rangle$ resonance, we have the following identities for their couplings:

$$
\begin{aligned}
g(a_0 \to \pi\eta_8) &= \frac{1}{\sqrt{5}}\, g_8 \ , \\
g(a_0 \to (K\bar{K})_1\rangle &= -\sqrt{\frac{3}{10}}\, g_8 \ , \\
g(\kappa \to (K\pi)_{1/2}) &= \frac{3}{\sqrt{20}}\, g_8 \ , \\
g(\kappa \to K\eta_8) &= -\frac{1}{\sqrt{20}}\, g_8 \ . \qquad (2.7)
\end{aligned}
$$

As a result, we predict the following three independent relations:

$$
\begin{aligned}
\frac{g(a_0 \to \pi\eta_8)}{g(a_0 \to (K\bar{K})_1)} &= -\sqrt{\frac{2}{3}} = -0.82 \ , \\
\frac{g(\kappa \to (K\pi)_{1/2})}{g(\kappa \to K\eta_8)} &= -3 \ , \\
\frac{g(a_0 \to (K\bar{K})_1)}{g(\kappa \to (K\pi)_{1/2})} &= -\sqrt{\frac{2}{3}} = -0.82 \ . \qquad (2.8)
\end{aligned}
$$

Averaging the values given in table 1 we obtain:

$$
\left|\frac{g(a_0 \to \pi\eta_8)}{g(a_0 \to (K\bar{K})_1)}\right| = 0.70 \pm 0.04 \ ,
$$

$$
\begin{aligned}
\left|\frac{g(\kappa\to(K\pi)_{1/2})}{g(\kappa\to K\eta_8)}\right| &= 1.64\pm 0.05,\\
\left|\frac{g(a_0\to(K\bar{K})_1)}{g(\kappa\to(K\pi)_{1/2})}\right| &= 1.10\pm 0.03\ .
\end{aligned}
\tag{2.9}
$$

For the first and third ratio the agreement is within the expected accuracy of around a 20%. It is worth stressing that the third ratio relates the couplings between both resonances and then is sensitive to the fact that g_8 is a common parameter in the SU(3) analysis. However, for the second ratio the disagreement is by a factor of two and hence for this particular coupling the SU(3) expectation does not hold within the expected accuracy. Nevertheless, we will show below that when SU(3) breaking is allowed in the SU(3) analysis we recover the agreement for the second ratio as well. Thus, these results, together with the observed soft evolution of the poles from the SU(3) limit to the physical one, support the picture of the $a_0(980)$ and the κ as members of the same octet. Still, we shall give below an additional analysis to support this conclusion. Notice that since the third ratio is in agreement with our analysis, the coupling $g(\kappa\to K\eta_8)$ is the one in disagreement with the present SU(3) expectations, as confirmed below when we allow SU(3) breaking linear in the quark masses.

In addition to the ratios in eq.(2.9) we can also obtain $|g_8|$ from the values of $|g(a_0\to\pi\eta_8)|$, $|g(a_0\to(K\bar{K})_1)|$ and $|g(\kappa\to(K\pi)_{1/2})|$ in table 1 and the SU(3) relations in eq.(2.7):

$$
|g_8| = 8.7\pm 1.3 \text{ GeV}\ ,
\tag{2.10}
$$

with a relative error around a 15%, well within the accuracy of an SU(3) analysis. In the number above we have not included the value of $|g(\kappa\to K\eta_8)|$ since, as discussed above, departs more than the allowed 20% from the SU(3) expectations at this stage.

Let us now consider the $I=0$ resonances, namely, the σ and the $f_0(980)$. These resonances are expressed as a mixture of the $I=0$ singlet and octet states, $|S_1\rangle$ and $|S_8\rangle$, respectively:

$$
\begin{aligned}
|\sigma\rangle &= \cos\theta|S_1\rangle+\sin\theta|S_8\rangle\ ,\\
|f_0(980)\rangle &= -\sin\theta|S_1\rangle+\cos\theta|S_8\rangle\ ,
\end{aligned}
\tag{2.11}
$$

with θ the mixing angle. Then we have the following relations for the couplings:

$$
\begin{aligned}
g(\sigma\to(\pi\pi)_0) &= -\frac{\sqrt{3}}{4}\cos\theta g_1-\sqrt{\frac{3}{10}}\sin\theta g_8\ ,\\
g(\sigma\to(K\bar{K})_0) &= -\frac{1}{2}\cos\theta g_1+\frac{1}{\sqrt{10}}\sin\theta g_8\ ,\\
g(\sigma\to(\eta_8\eta_8)_0) &= \frac{1}{4}\cos\theta g_1-\frac{1}{\sqrt{10}}\sin\theta g_8\ ,\\
g(f_0\to(\pi\pi)_0) &= \frac{\sqrt{3}}{4}\sin\theta g_1-\sqrt{\frac{3}{10}}\cos\theta g_8\ ,\\
g(f_0\to(K\bar{K})_0) &= \frac{1}{2}\sin\theta g_1+\frac{1}{\sqrt{10}}\cos\theta g_8\ ,\\
g(f_0\to(\eta_8\eta_8)_0) &= -\frac{1}{4}\sin\theta g_1-\frac{1}{\sqrt{10}}\cos\theta g_8\ .
\end{aligned}
\tag{2.12}
$$

From table 1 there is a very stark result, the almost vanishing coupling, much smaller than any other, of the σ resonance with the $\eta_8\eta_8$ channel. Imposing then that $g(\sigma \to \eta_8\eta_8) = 0$ in the equations above we obtain the relation:

$$\frac{g_1}{g_8} = \sqrt{\frac{8}{5}} \tan\theta \ . \tag{2.13}$$

Employing this result, it follows that:

$$\begin{aligned}
\frac{g(\sigma \to (\pi\pi)_0)}{g_8} &= -2\sqrt{\frac{3}{10}} \sin\theta \ , \\
\frac{g(f_0 \to (K\bar{K})_0)}{g_8} &= \frac{1}{\sqrt{10}} \cos\theta (1 + 2\tan^2\theta) \ .
\end{aligned}$$

We take the previous equations to calculate simultaneously g_8 and θ. Nevertheless, since we only know the modules of the different couplings, as shown in table 1, we can only determine the modules of θ and g_8. Notice that from eq.(2.13), the relative sign between g_1 and g_8 is fixed by the chosen one of θ. Once this is fixed, a change of sign in g_8 only gives rise to a global change of sign in the coupling constants listed in eq.(2.12) since from eq.(2.13) it implies a simultaneous change in the sign of g_1 as well. Thus, it is a matter of convention to choose the sign of g_8 and θ with our information of the modules of the coupling constants given in table 1. However, once the signs of g_8 and θ are chosen then from eq.(2.13) the sign of g_1 is fixed:

$$\begin{aligned}
|g_8| &= 10.5 \pm 0.5 \ \text{GeV} \ , \\
\cos^2\theta &= 0.934 \pm 0.006 \ , \\
|\theta| &= 14.9^o \pm 0.7^o \ .
\end{aligned} \tag{2.14}$$

Interestingly, the consistency of our approach is notorious by comparing the value of g_8 from the study of the a_0 and κ resonances with that in eq.(2.14) from the study of the mixing in the $I = 0$ system. We see that both values agree within errors which at most are a 15%. We show as well the average value of g_8 from the equation above and eq.(2.10), so that:

$$|g_8| = 9.6 \pm 1.3 \ \text{GeV}. \tag{2.15}$$

From eq.(2.13), we have the following value for g_1:

$$|g_1| = 3.5 \pm 0.5 \ \text{GeV} \ . \tag{2.16}$$

Now we use a more versatile way of treating eqs.(2.7) and (2.12) which is also more suited for the case when further SU(3) breaking is considered. For that we employ the method of the least squares. We construct the χ^2 function as:

$$\chi^2 = \sum_{R,PQ} \frac{|g(R \to PQ) - \hat{g}(R \to PQ)|^2}{\sigma^2_{R,PQ}} \ , \tag{2.17}$$

where the sum extends over all the resonances σ, $f_0(980)$, κ, $a_0(980)$ and all the pseudoscalar pairs PQ as shown in table 1. We have denoted by $\hat{g}(R \to PQ)$ those coupling constants calculated from the SU(3) analysis, while $g(R \to PQ)$ refers to the average ones in table 1.

In order to perform the numerical analysis based on the minimization of eq.(2.17), we have allowed for different signs in the evaluation of eqs.(2.12) since in table 1 we only know the modules of the coupling constants. The only consistent fit, where the ascribed signs for the couplings in the minimization process and those that finally come up coincide, is very similar to our previous results, eqs.(2.14), eq.(2.15) and (2.16). The precise values are:

$$\begin{aligned} |g_1| &= 3.9 \pm 0.8 \ \text{GeV} \ , \\ |g_8| &= 8.2 \pm 0.8 \ \text{GeV} \ , \\ |\theta| &= 19^o \pm 5^o \ , \end{aligned} \tag{2.18}$$

such that the relative sign of g_1 with respect to g_8 is in agreement with eq.(2.13). In order to calculate the errors we have multiplied by a common factor the errors of the coupling constants in the fifth column of table 1, much smaller than the expected accuracy of the SU(3) analysis, so that the χ^2 per degree of freedom is one. Our results are clearly compatible with the previously obtained values for g_8 (2.15), g_1 (2.16) and θ (2.14). The resulting couplings are:

$$\begin{aligned} |g(a_0 \to \pi\eta_8)| &= 3.7 \pm 0.4 \ \text{GeV} \ , \\ |g(a_0 \to (K\bar{K})_1)| &= 4.5 \pm 0.4 \ \text{GeV} \ , \\ |g(\kappa \to (K\pi)_{1/2}| &= 5.5 \pm 0.5 \ \text{GeV} \ , \\ |g(\kappa \to K\eta_8)| &= 1.8 \pm 0.2 \ \text{GeV} \ , \\ |g(\sigma \to (\pi\pi)_0)| &= 3.0 \pm 0.5 \ \text{GeV} \ , \\ |g(\sigma \to (K\bar{K})_0)| &= 0.9 \pm 0.5 \ \text{GeV} \ , \\ |g(\sigma \to (\eta_8\eta_8)_0)| &= 0.0 \pm 0.3 \ \text{GeV} \ , \\ |g(f_0 \to (\pi\pi)_0)| &= 3.7 \pm 0.5 \ \text{GeV} \ , \\ |g(f_0 \to (K\bar{K})_0)| &= 3.1 \pm 0.3 \ \text{GeV} \ , \\ |g(f_0 \to (\eta_8\eta_8)_0)| &= 2.8 \pm 0.2 \ \text{GeV} \ . \end{aligned} \tag{2.19}$$

We see a remarkable agreement between the calculated couplings from our SU(3) analysis in eq.(2.19) and the averages reported in the fifth column of table 1. Of the ten couplings there are only two exceptions. The already discussed $g(\kappa \to K\eta_8)$ which turns around a factor 2 too small and the $g(f_0 \to (\pi\pi)_0)$ which is too large by a factor around 3. It should be noted that the $|g(f_0 \to (\pi\pi)_0)|$ coupling is the most sensitive to the presence or absence of a preexisting state in the $f_0(980)$ resonance since it changes by a 40%, while for the other couplings the change is lower than a 5%. This points towards a special status for this coupling which controls the width of the $f_0(980)$ resonance. Indeed the large reduction, by a factor of 2, between the imaginary part of the pole position of the $f_0(980)$ resonance in the third column of table 1 reflects the reduction in the value of its coupling to the $(\pi\pi)_0$ system. Notice that the ratio of the square of the couplings of the f_0 to $|\pi\pi\rangle_0$ given in the second and third columns is $(0.89/1.33)^2 \simeq 1/2$, as corresponds to the ratio of the widths. In addition, as noticed in ref.[32, 35], the $f_0(980)$ behaves as a pure $K\bar{K}$

Resonance	Full ref.[22] GeV	No bare poles eq.(2.1) GeV	No bare poles eq.(2.2) GeV	Average
σ	$\lvert g(\sigma\to 1)\rvert = 3.65$ $\lvert g(\sigma\to 8_s)\rvert = 2.67$	$\lvert g(\sigma\to 1)\rvert = 3.86$ $\lvert g(\sigma\to 8_s)\rvert = 2.38$	$\lvert g(\sigma\to 1)\rvert = 3.89$ $\lvert g(\sigma\to 8_s)\rvert = 2.38$	$\lvert g(\sigma\to 1)\rvert = 3.80\pm 0.13$ $\lvert g(\sigma\to 8_s)\rvert = 2.48\pm 0.18$
$f_0(980)$	$\lvert g(f_0\to 1)\rvert = 5.35$ $\lvert g(f_0\to 8_s)\rvert = 4.15$	$\lvert g(f_0\to 1)\rvert = 5.07$ $\lvert g(f_0\to 8_s)\rvert = 3.90$	$\lvert g(f_0\to 1)\rvert = 4.20$ $\lvert g(f_0\to 8_s)\rvert = 2.45$	$\lvert g(f_0\to 1)\rvert = 4.9\pm 0.6$ $\lvert g(f_0\to 8_s)\rvert = 3.5\pm 0.9$
a_0	$\lvert g_8\rvert = 8.95$	$\lvert g_8\rvert = 8.83$	$\lvert g_8\rvert = 9.01$	$\lvert g_8\rvert = 8.5\pm 0.5$
κ	$\lvert g_8\rvert = 8.12$	$\lvert g_8\rvert = 7.94$	$\lvert g_8\rvert = 7.90$	$\lvert g_8\rvert = 7.99\pm 0.12$

Table 2: Coupling constants of the scalar resonances with two mesons states with well defined SU(3) eigenvalues. In the fifth column we show the average of the values of the three previous columns.

bound state when the $\pi\pi$ channel is removed. As a result, one expects that the $g(f_0 \to (\pi\pi)_0)$ coupling is mostly driven by kaonic loops decaying into two pions so that the SU(3) breaking can be huge due to the large difference between the $K\bar{K}$ and $\pi\pi$ thresholds.

Summarizing, of the ten couplings constant we have used three of them to calculate θ, g_8, g_1, and then we end with the values given in eq.(2.19). Between the latter, eight couplings are in very good agreement with their 'experimental' values given in table 1 but two of them are in disagreement, although the coupling $g(\kappa \to K\eta_8)$ will agree with our SU(3) analysis once further SU(3) breaking corrections are taken into account, eq.(2.24). On the other hand, we have also checked that the obtained value of g_8 from the study of the pure octet states κ and $a_0(980)$, eq.(2.10), is compatible with that from the study of the mixing in the $I = 0$ system, eq.(2.14). The average of both values is the one quoted in eq.(2.15), compatible with the value of g_8 given in eq.(2.18) from the minimization of the χ^2 function, eq.(2.17). This is an important check and then, as a whole picture, we can claim that the σ, $f_0(980)$, $a_0(980)$ and κ constitute a nonet of dynamically generated scalar resonances with a mixing angle of around 20^o. The preexisting bare pole contributing to the $f_0(980)$ resonance appears as a small contribution to the $f_0(980)$ physical pole that particularly affects the couplings of this resonance with the $\pi\pi$ channel, as apparent when comparing the couplings of the $f_0(980)$ in the second column of table 1 with those in the third one.

As stated above, we consider a second SU(3) coupling constant analysis to support our view that these resonances constitute a nonet. We now consider the couplings shown in table 2 of the σ, $f_0(980)$, $a_0(980)$ and κ scalar resonances with the two pseudoscalar states of eq.(2.5) with well defined SU(3) eigenvalues.

Since the $a_0(980)$ and κ are pure octet resonances, their couplings to an octet state must be the same and this is indeed what happens as can be seen from the values given in table 2, which are the same within an error of around 9%. Averaging the values of g_8 in the first three columns of table 2, we have:

$$|g_8| = 8.5 \pm 0.5 \text{ GeV} , \tag{2.20}$$

a very similar value to that of eq.(2.18). The weighted value of g_8 between the ones given in eqs.(2.15) and (2.20) is:

$$|g_8| = 8.6 \pm 0.5 \text{ GeV} \ . \tag{2.21}$$

Because the f_0 and σ are a mixture between the pure singlet and the $I=0$ octet states, eq.(2.11), their couplings to the singlet and octet states are driven by this mixing and then we have:

$$\begin{aligned} g(\sigma \to 1) &= \cos\theta \, g_1 \ , \\ g(\sigma \to 8_s) &= \sin\theta \, g_8 \ , \\ g(f_0 \to 1) &= -\sin\theta \, g_1 \ , \\ g(f_0 \to 8_s) &= \cos\theta \, g_8 \ . \end{aligned} \tag{2.22}$$

Taking into account the second line of the previous equations and the calculated value of g_8 in eq.(2.21), we then have a new calculation for g_1 and the mixing angle θ.

$$\begin{aligned} |g_1| &= 3.96 \pm 0.13 \text{ GeV} \ , \\ \cos^2\theta &= 0.916 \pm 0.006 \ , \\ |\theta| &= 16.8^o \pm 0.7^o \ , \end{aligned} \tag{2.23}$$

very similar to the values of eq.(2.18) and compatible within the shown errors from the dispersion in the values shown in tables 1 and 2.

Let us notice that we can also find a solution for the last two equations in (2.22). One then obtains very different values to those of eqs.(2.18) and (2.23) with $\theta \simeq 65^0$, for g_8 given in eq.(2.20). Indeed it is easy to see by finding $\tan\theta$ from either of the ratios $g(\sigma \to 8_s)/g(f_0 \to 8_s)$ or $g(f_0 \to 1)/g(\sigma \to 1)$ that the set of four equations in (2.22) is not compatible with the values given in table 2. This is why we have solved separately those two equations of the σ, whose solution is given in eq.(2.23), and those of the $f_0(980)$. Indeed, the χ^2 (2.17) calculated from the set of values of eq.(2.23) is 1.5 smaller than that calculated from the values that result if we considered instead the f_0 couplings in eqs.(2.22). Furthermore, if we take the latter as starting values to minimize the χ^2 function, we end then with a solution similar to eqs.(2.18) and (2.23). Let us note from eq.(2.5) that the two pion system enters both in the octet and singlet $I=0$ states with large SU(3) Clebsch-Gordan coefficients and then this last method cannot be applied to the $f_0(980)$ since it was not good enough for the study of the $g(f_0 \to (\pi\pi)_0)$ coupling. It is also worth stressing that the problem with the $g(f_0 \to (\pi\pi)_0)$ coupling cannot be solved by considering an extra singlet contribution to the $f_0(980)$ resonance, as one could think because of the preexisting singlet contribution found in ref.[22] to the physical $f_0(980)$. This is so because the $g(f_0 \to 8_s)$ coupling does not involve any singlet contribution and is in disagreement with our expectations from eq.(2.22), with the values determined in eqs.(2.20) and (2.23) from the sudy of the couplings of the σ, $a_0(980)$ and κ resonances in table 2.

As state above we now consider SU(3) breaking in the couplings of the octet and singlet states to two pseudoscalars. Since the singlet and the symmetric octet SU(3) representations are symmetric under the exchange of the two pseudoscalars, at linear order in the quark mass matrix, we can write the following Lagrangian:

$$F\langle S(2\Phi\chi\Phi + \Phi^2\chi + \chi\Phi^2)\rangle + G S_1 \langle 2\Phi\chi\Phi + \Phi^2\chi + \chi\Phi^2\rangle \ . \tag{2.24}$$

The same expression can be deduced from the chiral Lagrangians of ref.[47]. In the expression above Φ is the usual 8×8 matrix with the octet of pseudoscalars and the same applies for S but now referring to the scalar octet. For more details see e.g. ref.[47]. Finally, S_1 is the scalar singlet, $\chi = diag(0,0,m_K^2 - m_\pi^2) \propto (0,0,m_s - \hat{m})$ is the matrix responsible for the SU(3) breaking with m_s the strange quark mass and $\hat{m}$ the average of the masses of the two lightest quarks. On the other hand, F and G are two constants. We have then minimized the χ^2, eq.(2.17), with the corrections derived from the previous Lagrangian added to eqs.(2.7) and (2.12). We want to emphasize here that the values of the couplings g_8, g_1 and θ turn out to be the same and the SU(3) breaking parameters $F(m_K^2 - m_\pi^2) = -0.23 \pm 0.16$ and $G(m_K^2 - m_\pi^2) = 0.13 \pm 0.4$ take very natural values, around a 20% of unity as expected from SU(3) breaking. The main novelty is that now $g(\kappa \to K\eta_8) = 2.52$, quite close to the value given in table 1. Nevertheless, the situation for the $g(f_0 \to (\pi\pi)_0)$ coupling does not improve and we are still a factor 3 off. For completeness, we also discuss here a qualitative different fit to that of eq.(2.18) where the SU(3) breaking is much larger but the χ^2 function is smaller, by around a factor of 2, than that corresponding to the fit in eq.(2.18) but including now the just mentioned SU(3) breaking corrections. This fit arises when allowing a change of sign in the coupling $g(\kappa \to K\eta_8)$ with respect to the case without the SU(3) breaking from eq.(2.24). Let us note that this coupling is SU(3) suppressed because of the smallness of the corresponding Clebsch-Gordan, see eq.(2.5), so that moderate SU(3) corrections could well change its sign. However, in this case the SU(3) breaking parameters are larger by around a factor of 7 to those given above, so that $F(m_K^2 - m_\pi^2) = 1.44 \pm 0.11$ and $G(m_K^2 - 6m_\pi^2) = 0.81 \pm 0.2$ and hence they drive to large SU(3) corrections that are difficult to accept. We see e.g. that the resulting g_8 parameter is lower by a factor of two than that obtained directly from the $a_0(980)$ and κ analysis in eq.(2.20), by directly averaging the values shown in table 2. This is of course very unlikely to be accepted since we see a nice agreement between the couplings of the $a_0(980)$ and κ to the pure octet states in table 2 as corresponds to a soft SU(3) breaking. Furthermore, from this fit to the two pseudoscalar couplings we can then predict the values of the couplings of the resonances to the SU(3) eigenstates in eq.(2.5) and then we observe that the predicted value for g_8 is much smaller, by around a factor 20, than the ones shown in table 2. As a result we discard this fit as non-physical and keep the one of eq.(2.18) as our final results.

By passing, we mention that taking the real part of the poles presented in table 1 as the masses of the corresponding resonances, we then find that they fulfill in good agreement a *linear* Gell-Mann-Okubo mass relation with the mixing angle given in eq.(2.18):

$$4m_\kappa - m_{a_0} = 3(m_{f_0}\cos^2\theta + m_\sigma \sin^2\theta) \ . \tag{2.25}$$

The ratio between the difference of the right(2.7) and left(2.1) sides of the above equation and the average of both is around a 25%, well shaped within the usual SU(3) accurateness. On the other hand, the quadratic Gell-Mann-Okubo mass relation is much worse behaved, being the relative difference around a factor 2.

Finally, the sign of θ can be fixed by considering the strength of the scalar form factors:

$$\begin{aligned} f_n &= \langle 0|\frac{1}{\sqrt{2}}(\bar{u}u + \bar{d}d)|K\bar{K}\rangle_0 \ , \\ f_s &= \langle 0|\bar{s}s|K\bar{K}\rangle_0 \ , \end{aligned}$$

$$(2.26)$$

at the peak of the $f_0(980)$. In the previous equations we have denoted by $|0\rangle$ the vacuum state. These form factors, among others, were calculated in ref.[48], and the peak of the $f_0(980)$ is clearly visible. Now, from ref.[48] at the $f_0(980)$ peak the ratio $|f_s/f_n| \simeq 5.5$, denoting clearly that the $f_0(980)$ couples stronger with strangeness, as well known. Notice that the coupling of the $f_0(980)$ to the two kaons factorizes out in the ratio and then one is only sensitive to the f_0 coupling to the quark-antiquark sources. We can perform the SU(3) decomposition of the previous sources:

$$\begin{aligned}\bar{s}s &= \frac{1}{\sqrt{3}}s_1 - \sqrt{\frac{2}{3}}s_8 \ , \\ \bar{n}n \equiv \frac{1}{\sqrt{2}}(\bar{u}u + \bar{d}d) &= \sqrt{\frac{2}{3}}s_1 + \frac{1}{\sqrt{3}}s_8 \ , \end{aligned} \tag{2.27}$$

where s_1 and s_8 are singlet and octet operators. Then, neglecting any other $SU(3)$ breaking beyond mixing we can write:

$$\begin{aligned}\langle 0|\bar{s}s|f_0(980)\rangle &= -\sin\theta\frac{1}{\sqrt{3}}\langle 0|s_1|S_1\rangle - \cos\theta\sqrt{\frac{2}{3}}\langle 0|s_8|S_8\rangle \ , \\ \langle 0|\bar{n}n|f_0(980)\rangle &= -\sin\theta\sqrt{\frac{2}{3}}\langle 0|s_1|S_1\rangle + \cos\theta\frac{1}{\sqrt{3}}\langle 0|s_8|S_8\rangle \ . \end{aligned} \tag{2.28}$$

Now, from the value $|\theta| = 19^o$ given in eq.(2.18) and taking, just for an order of magnitude estimation, $\langle 0|s_1|S_1\rangle \simeq \langle 0|s_8|S_8\rangle$, as required by U(3) symmetry, we then have:

$$\begin{aligned}\theta = +19^o \quad &\rightarrow \quad \left|\frac{f_s}{f_n}\right| = 3.5 \ , \\ \theta = -19^o \quad &\rightarrow \quad \left|\frac{f_s}{f_n}\right| = 0.7 \ . \end{aligned} \tag{2.29}$$

Thus, the solution with $\theta > 0$ is clearly favored since requires much smaller U(3) breaking corrections than that with $\theta < 0$, which would be huge. As a result, if we take the convention that $g_8 > 0$, we can remove the absolute value signs in eq.(2.18), and write:

$$\begin{aligned} g_8 &= 8.2 \pm 0.8 \ \text{GeV} \ , \\ g_1 &= 3.9 \pm 0.8 \ \text{GeV} \ , \\ \theta &= +19^o \pm 5^o \end{aligned} \tag{2.30}$$

It is also worth mention that our mixing angle is in agreement with the determination from ref.[11].

3 Conclusions

We have first shown that the σ, $f_0(980)$, $a_0(980)$ and κ poles evolve continuously in the λ parameter, eq.(2.3), from the physical limit $\lambda = 0$ to a nonet of resonances in a SU(3) limit case,

$\lambda = 1$, made up by a degenerate octet and a singlet. Indeed we already showed this in ref.[22], but now we have analysed in further detail the situation where the bare poles, an octet with a mass around 1.4 GeV and a singlet with a mass around 1 GeV, are removed. In particular, we have now performed two fits to the scattering data of the S-waves in the $I = 0$, 1/2 and 1 channels without bare resonances and with a common value for all the subtractions constants, eq.(2.1), or with different ones and without the $\eta_8\eta_8$ channel as well, eq.(2.2). In addition, we have also interpreted that a value around -1 for the subtraction constants at the scale $\mu = M_\rho$ corresponds to calculating the unitarity loops with a three-momentum cut-off of a natural value, $Q \simeq M_\rho$. Although this certainly shows that the aforementioned resonances constitute a nonet of dynamically generated scalar resonances with the $a_0(980)$ and κ being members only of the octet, we have performed two additional analyses that have shown: First, the SU(3) limit picture as a whole is still useful in order to study the lightest scalar nonet in the physical limit (within the expected precision of around a 20% for such estimations). Second, we consistently conclude again that those resonances belong to the same nonet and third, we are able to obtain a value for the mixing angle between the $I = 0$ states of the singlet and octet multiplets, together with the coupling constants g_8 and g_1. In this way we consistently study within SU(3) symmetry, in terms of three parameters, the set of couplings: $g(a_0 \rightarrow (K\bar{K})_1)$, $g(a_0 \rightarrow \pi\eta_8)$, $g(a_0 \rightarrow 8_s)$, $g(\kappa \rightarrow (K\pi)_{1/2})$, $g(\kappa \rightarrow K\eta_8)$, $g(\kappa \rightarrow 8_s)$, $g(\sigma \rightarrow (\pi\pi)_0)$, $g(\sigma \rightarrow (K\bar{K})_0)$, $g(\sigma \rightarrow (\eta_8\eta_8)_0)$, $g(\sigma \rightarrow 1)$, $g(\sigma \rightarrow 8_s)$, $g(f_0 \rightarrow (K\bar{K})_0)$, $g(f_0 \rightarrow (\eta_8\eta_8)_0)$. We have seen that the coupling $g(\kappa \rightarrow K\eta_8)$ is finally in agreement with our SU(3) analysis once linear SU(3) breaking in the quark masses is taken into account from eq.(2.24). Nevertheless, the coupling $g(f_0(980) \rightarrow (\pi\pi)_0)$ require still higher orders in the SU(3) breaking parameters and do not follow the standard simple scheme developed here in the lines of the well established vector and tensor nonets. The given SU(3) analyses take the values of the couplings of the scalar resonances with two pseudoscalar states with either well defined isospin or well defined SU(3) eigenvalues. Both analyses have different sensitivity to the SU(3) breaking, since different combinations of amplitudes are involved, and follow very different numerical analyses, but remarkably, compatible values between errors for the mixing angle as well as for the coupling g_8 and g_1 stem. This consistency is worth stressing and gives us further confidence in our results together with our interpretation that the failure in reproducing the $(\pi\pi)_0$ coupling of the $f_0(980)$ should be considered as an isolated fact in an overall good description and interpretation of table 1. We have also established that the scalar resonances fulfill to good precision a linear Gell-Mann-Okubo mass relation with our determined mixing angle and masses in table 1. Finally, from a study of the scalar form factors calculated in ref.[48], we have fixed the sign of the mixing angle to be positive and then we conclude that $\theta = +19^o \pm 5^o$.

Acknowledgments

I would like to acknowledge useful discussions with D. Black and E. Torrente. This work is partially supported by the DGICYT project FPA2002-03265.

References

[1] S. D. Protopopescu *et al.*, Phys. Rev. D **7** (1973) 1279.

[2] R. Ammar *et al.*, Phys. Rev. Lett. **21** (1968) 1832.

[3] D. Morgan, Phys. Lett. B **51** (1974) 71.

[4] J. J. Sakurai, Phys. Rev. Lett. **9** (1962) 472.

[5] S. Okubo, Phys. Lett. **5** (193) 165.

[6] S. L. Glashow and R. H. Socolow, Phys. Rev. Lett. **15** (1965) 329.

[7] V. Cirigliano, G. Ecker, H. Neufeld and A. Pich, hep-ph/0305311.

[8] R. L. Jaffe, Phys. Rev. D **15**, 267 (1977).

[9] A. Chodos, R. L. Jaffe, K. Johnson, C. B. Thorn and V. F. Weisskopf, Phys. Rev. D **9** (1974) 3471.

[10] M. D. Scadron, Phys. Rev. D **26** (1982) 239.

[11] M. Napsuciale, arXiv:hep-ph/9803396.

[12] N. Tornqvist, Eur. Phys. J. **C11** (1999) 359.

[13] G.'t Hooft, hep-th/9903189.

[14] J. Schechter and Y. Ueda, Phys. Rev. D **3** (1971) 2874.

[15] M. Levy, Nuov. Cim. LIIA (1967) 7247.

[16] G. t'Hooft, Phys. Rep. 142 (1986) 357.

[17] D. Black, A. H. Fariborz, F. Sannino and J. Schechter, Phys. Rev. D **59** (1999) 074026 [arXiv:hep-ph/9808415].

[18] E. van Beveren and G. Rupp, arXiv:hep-ph/0304105.
E. Van Beveren, T. A. Rijken, K. Metzger, C. Dullemond, G. Rupp and J. E. Ribeiro, Z. Phys. C **30**, 615 (1986).

[19] M. Ishida, Prog. Theor. Phys. **101** (1999) 661 [arXiv:hep-ph/9902260].

[20] J. A. Oller, E. Oset and J. R. Pelaez, Phys. Rev. D **59** (1999) 074001 [Erratum-ibid. D **60** (1999) 099906] [arXiv:hep-ph/9804209].

[21] A. Gomez Nicola and J. R. Pelaez, Phys. Rev. D **65** (2002) 054009 [arXiv:hep-ph/0109056].
J. R. Pelaez and A. Gomez Nicola, AIP Conference Proceedings 660, pag. 102. Editors, A. H. Blin et al. arXiv:hep-ph/0301049.

[22] J. A. Oller and E. Oset, Phys. Rev. D **60** (1999) 074023 [arXiv:hep-ph/9809337].

[23] M. Jamin, J. A. Oller and A. Pich, Nucl. Phys. B **587** (2000) 331 [arXiv:hep-ph/0006045].

[24] G. Janssen, B. C. Pearce, K. Holinde and J. Speth, Phys. Rev. D **52** (1995) 2690 [arXiv:nucl-th/9411021].
D. Lohse, J. W. Durso, K. Holinde and J. Speth, Nucl. Phys. A **516** (1990) 513.

[25] F. E. Close and N. A. Tornqvist, J. Phys. G **28** (2002) R249 [arXiv:hep-ph/0204205].

[26] K. Hagiwara *et al.* [Particle Data Group Collaboration], Phys. Rev. D **66** (2002) 010001.

[27] A. Aloisio *et al.* [KLOE Collaboration], Phys. Lett. B **537** (2002) 21 [arXiv:hep-ex/0204013].
A. Aloisio *et al.* [KLOE Collaboration], Phys. Lett. B **536** (2002) 209 [arXiv:hep-ex/0204012].

[28] J. A. Oller, Nucl. Phys. A **714** (2003) 161 [arXiv:hep-ph/0205121].

[29] A. Bramon, R. Escribano, J. L. Lucio M., M. Napsuciale and G. Pancheri, Phys. Lett. B **494** (2000) 221 [arXiv:hep-ph/0008188].
A. Bramon, R. Escribano, J. L. Lucio M, M. Napsuciale and G. Pancheri, Eur. Phys. J. C **26** (2002) 253 [arXiv:hep-ph/0204339].

[30] N. N. Achasov and A. V. Kiselev, arXiv:hep-ph/0212153.

[31] M. Boglione and M. R. Pennington, arXiv:hep-ph/0303200.

[32] F. E. Close, N. Isgur and S. Kumano, Nucl. Phys. B **389** (1993) 513 [arXiv:hep-ph/9301253].

[33] E. M. Aitala *et al.* [E791 Collaboration], Phys. Rev. Lett. **86** (2001) 765 [arXiv:hep-ex/0007027]; Phys. Rev. Lett. **86** (2001) 770 [arXiv:hep-ex/0007028]; Phys. Rev. Lett. **89** (2001) 121801.

[34] C. Göbel, arXiv:hep-ex/0012009.

[35] J. A. Oller and E. Oset, Nucl. Phys. A **620** (1997) 438 [Erratum-ibid. A **652** (1999) 407] [arXiv:hep-ph/9702314].
For a general introduction and discussion see J. A. Oller, proceeding of the YITP Workshop on Possible Existence on the Sigma Meson and its Implications to Hadron Physics, Soryushiron Kenkyu (Kyoto) 102, No. 5 (2001) 33-40, [arXiv:hep-ph/0007349].

[36] J. A. Oller and E. Oset, Nucl. Phys. A **629** (1998) 739 [arXiv:hep-ph/9706487].

[37] U. G. Meißner and J. A. Oller, Nucl. Phys. A **679** (2001) 671 [arXiv:hep-ph/0005253].

[38] J. A. Oller, Phys. Lett. B **426** (1998) 7 [arXiv:hep-ph/9803214].

[39] S. Weinberg, Physica **A96**, 327 (1979).

[40] J. Gasser and H. Leutwyler, Ann. Phys. (N.Y.) **158**, 142 (1984).

[41] J. A. Oller, arXiv:nucl-th/0207086, to be published in Nucl. Phys. A; J. A. Oller and U. G. Meißner, Phys. Lett. B **500** (2001) 263 [arXiv:hep-ph/0011146]; U. G. Meißner and J. A. Oller, Nucl. Phys. A **673** (2000) 311 [arXiv:nucl-th/9912026];

[42] D. Jido, J. A. Oller, E. Oset, A. Ramos and U. G. Meissner, arXiv:nucl-th/0303062.

[43] K. L. Au, D. Morgan and M. R. Pennington, Phys. Rev. D **35** (1987) 1633.

[44] P. Estabrooks, R. K. Carnegie, A. D. Martin, W. M. Dunwoodie, T. A. Lasinski and D. W. Leith, Nucl. Phys. B **133** (1978) 490.

[45] D. Aston *et al.*, Nucl. Phys. B **296** (1988) 493.

[46] M. Jamin, J. A. Oller and A. Pich, Nucl. Phys. B **622** (2002) 279 [arXiv:hep-ph/0110193].

[47] G. Ecker, J. Gasser, A. Pich and E. de Rafael, Nucl. Phys. B **321**, 311 (1989).

[48] U. G. Meissner and J. A. Oller, Nucl. Phys. A **679**, 671 (2001) [arXiv:hep-ph/0005253].